

ON CLOSED UNBOUNDED SETS CONSISTING
OF FORMER REGULARS

Moti Gitik

School of Mathematical Sciences
Sackler Faculty of Exact Sciences
Tel Aviv University
Ramat-Aviv, 69978 Israel

Abstract

A method of iteration of Prikry type forcing notions as well as a forcing for adding clubs is presented. It is applied to construct a model with a measurable cardinal containing a club of former regulars, starting with $o(\kappa) = \kappa + 1$. On the other hand, it is shown that the strength of above is at least $o(\kappa) = \kappa$.

Suppose that κ is an inaccessible cardinal. We wish to find a generic extension (usually cardinal preserving) such that $\{\alpha < \kappa \mid \alpha \text{ is regular in } V\}$ contains a club. Radin [Ra] introduced a basic method to do this. Simply start with a measurable κ with $o(\kappa) = \kappa^+$ and then force with the Radin forcing constructed from $o(\kappa) = \kappa^+$. If one wishes to keep κ a measurable in the extension, then a weak repeat point suffices. Both facts are proved by Mitchell [Mi1],

We show how to reduce assumptions rendering the above possible.

A method of iteration generalizing those of [Gi1] is presented. Then a variant of it is used to iterate forcing for shooting clubs. We think that this method of iteration can be applied to other distributive forcings as well.

We like to thank the referee for pointing out that the proof of section 2 gives only $o(\kappa) = \kappa$ and not $o(\kappa)\kappa + 1$ as was claimed in the previous version, for long and detailed list of corrections and for his requests on explaining certain parts of the paper.

1. A Forcing Construction

We will now prove the following two theorems.

Theorem 1.1. *Suppose that κ is an inaccessible cardinal such that for every $\delta < \kappa$ the set of α 's below κ with $o(\alpha) \geq \delta$ is stationary. Then there is a cardinal preserving generic extension such that the set $\{\alpha < \kappa \mid \alpha \text{ is regular in } V\}$ contains a club.*

Theorem 1.2. *Suppose that κ is a measurable cardinal with $o(\kappa) = \kappa + 1$. Then there is a cardinal preserving extension satisfying the following:*

- (1) κ is a measurable,
- (2) $\{\alpha < \kappa \mid \alpha \text{ is a regular in } V \text{ contains a club}\}.$

The proofs of these theorems use an iteration Prikry type forcing notion that was introduced in [Gi1].

Basically for every $\alpha < \kappa$ with $\alpha > o(\alpha) > o$ we are forcing a Prikry or Magidor sequence to α without adding new bounded subsets. The order type of the sequence is $\omega^{o(\alpha)}$, where the exponentiation is the ordinal one. The forcing used for this is $\langle \mathcal{P}(\alpha, o(\alpha)), \leq, \leq^* \rangle$

which was introduced in [Gi1]. The definition of it is quite long and will not be used explicitly here. The only facts used in connection with this are stated below.

Fact 1. $\langle \mathcal{P}(\alpha, o(\alpha)), \leq, \leq^* \rangle$ satisfies Prikry condition and it is α -weakly closed, i.e.

- (a) $\leq \supseteq \leq^*$;
- (b) for every $p \in \mathcal{P}(\alpha, o(\alpha))$ and every statement σ of the forcing $\langle \mathcal{P}(\alpha, o(\alpha)), \leq \rangle$ there is $q \geq^* p$ deciding σ ; (c) $\leq^*$ is α -closed.

Fact 2. $\langle \mathcal{P}(\alpha, o(\alpha)), \leq \rangle$ satisfies α^+ -c.c. and has cardinality 2^α .

Fact 3. $\langle \mathcal{P}(\alpha, o(\alpha)), \leq \rangle$ adds a closed cofinal in α sequence consisting of regular in V cardinals of order type $\omega^{o(\alpha)}$ and almost containing in every club of α of V .

We are going to iterate $\mathcal{P}(\alpha, 0(\alpha))$'s using the iteration of [Gi1]. For the benefit of the reader let us give a precise definition.

Let A be a set consisting of α 's such that $\alpha < \kappa$ and $\alpha > o(\alpha) > 0$. Denote by A^ℓ the closure of the set $\{\alpha + 1 \mid \alpha \in A\} \cup A$. For every $\alpha \in A^\ell$ define by induction $\mathcal{P}_\alpha$ to be the set of all elements p of the form $\langle p_\gamma \mid \gamma \in g \rangle$ where

- (1) g is a subset of $\alpha \cap A$.
- (2) g has an Easton support, i.e. for every inaccessible $\beta \leq \alpha$, $\beta > |\text{dom } g \cap \beta|$;
- (3) $p \restriction \gamma = \langle p_\beta \mid \beta \in \gamma \cap g \rangle \in \mathcal{P}_\gamma$ and $p \restriction \gamma \Vdash_{\mathcal{P}_\gamma} "p_\gamma \in \mathcal{P}(\gamma, o(\gamma))"$ for every $\gamma \in \text{dom } g$

Let $p = \langle p_\gamma \mid \gamma \in g \rangle$, $q = \langle q_\gamma \mid \gamma \in f \rangle$ be elements of $\mathcal{P}_\alpha$. Then $p \geq q$ (p is stronger than q) if the following holds:

- (1) $g \supseteq f$
- (2) for every $\gamma \in f$ $p \restriction \gamma \Vdash_{\mathcal{P}_\gamma} "p_\gamma \geq q_\gamma$ in the forcing $\mathcal{P}(\gamma, 0(\gamma))"$
- (3) there exists a finite subset b of f so that for every $\gamma \in f \setminus b$, $p \restriction \gamma \Vdash "p_\gamma \geq^* q_\gamma$ in $\mathcal{P}(\gamma, 0(\gamma))"$.

If $b = \emptyset$, then let $p \geq^* q$.

By [Gi1], such iteration preserves cardinals. The final forcing $\mathcal{P}_\kappa$ satisfies κ - c.c. We refer to [Gi1] for more details.

Proof of Theorem 1. Let G_κ be a generic subset of $\mathcal{P}_\kappa$. We force over $V[G_\kappa]$ with the forcing $P[E] = \{d \mid d \text{ is closed, bounded subset of } E\}$ ordered by the end extension, where

$$E = \{\alpha < \kappa \mid \alpha \text{ is a regular in } V\},$$

i.e. the usual forcing for adding a club via a stationary set.

By Avraham-Shelah [Av-Sh], if E is fat then this forcing is (κ, ∞) -distributive. Where fatness means that for every $\delta < \kappa$ and every closed unbounded subset C of κ there is a closed subset $s \subseteq C \cap E$ of order type δ .

So it is enough to show that E is fat in $V[G_\kappa]$. Let $\delta < \kappa$ and $C \subseteq \kappa$ be a club in $V[G_\kappa]$. By κ -c.c. of the forcing $\mathcal{P}_\kappa$ there is $C^* \subseteq C$ a club which belongs to V . Then pick an α^* a limit point of $C^* \cap \{\alpha < \kappa \mid o(\alpha) \geq \delta\}$. By Fact 3. there will be a closed subset of $C^* \cap \alpha^* \cap E$ of order type δ . This completes the proof. $\square$

We now turn to the proof of Theorem 1.2 which takes up the rest of this section.

Proof of Theorem 1.2. Denote by U the measure concentrating on α 's with $o(\alpha) = \alpha$.

We preserve the notation of Theorem 1.1.

Let us explain the idea of the proof. We like to force with $P[E]$ a club through the set E of regular cardinals below κ . So lots of changes of cofinalities are needed. An additional task is to preserve measurability of κ . So for a set of α 's in U we need to use $P[E \cap \alpha]$. The problem with it is that this forcing is not closed or weakly closed but only distributive. Iteration of $P[E \cap \alpha]$'s in Easton fashion even ω -times collapses cardinals. The key idea will be to embed $P[E \cap \alpha]$ into $\mathcal{P}(\alpha, \beta) * \beta$ -closed, where $\beta < \alpha$ and we will allow to change β from time to time in order to increase the degree of closeness. Then the scheme of iteration of [Gi1] will be generalized to make possible the iteration of less closed forcing notions.

Let us start with a generalization of the iteration process of [Gi1].

We describe a general scheme of iteration. A specific forcing used for the proof of 1.2 will be defined later.

Let A be a set consisting of inaccessible cardinals. Denote by A^ℓ the closure of the set $A \cup \{\alpha + 1 \mid \alpha \in A\}$. We define an iteration $\langle \mathcal{P}_\alpha, \mathcal{Q}_\alpha \mid \alpha \in A^\ell \rangle$. For every $\alpha \in A^\ell$ define by induction $\mathcal{P}_\alpha$ to be the set of all elements p of the form $\langle p_\gamma \mid \gamma \in g \rangle$, where

- (1) g is a subset of $\alpha \cap A$;
- (2) g has an Easton support, i.e. for every inaccessible $\beta \leq \alpha$ $\beta > |\text{dom } g \cap \beta|$;

(3) for every $\gamma \in \text{dom } g$

$$p \restriction \gamma = \langle p_\beta \mid \beta < \gamma \rangle \in \mathcal{P}_\gamma$$

and $p \restriction \gamma \Vdash_{\mathcal{P}_\gamma}$ “ p_γ is a condition in the forcing $\langle Q_\gamma, \leq_\gamma, \leq_\gamma^* \rangle$ satisfying the Prikry condition and of cardinality below the next element of $\check{A}$ above $\check{\gamma}$ ”.

The difference here is that we do not require that $\langle Q_\gamma, \leq_\gamma^* \rangle$ is γ -closed.

Let $p = \langle p_\gamma \mid \gamma \in g \rangle$ and $q = \langle q_\gamma \mid \gamma \in f \rangle$ be elements of $\mathcal{P}_\alpha$. Then $p \geq q$ (p is stronger than q) if the following holds:

(1) $g \supseteq f$

(2) for every $\gamma \in f$

$$p \restriction \gamma \Vdash_{\mathcal{P}_\gamma} “p_\gamma \geq_\gamma q_\gamma \text{ in the forcing } Q_\gamma”$$

(3) there exists a finite subset b of f so that for every $\gamma \in f \setminus b$ $p \restriction \gamma \Vdash_{\mathcal{P}_\gamma}$ “ $p_\gamma \geq_\gamma^* q_\gamma$ in the forcing Q_γ ”.

If the set b in (3) is empty we call p a direct extension of q and denote this by $p \geq^* q$.

Lemma 1.3. *Let $\alpha \in A^\ell$, $p \in \mathcal{P}_\alpha$ and σ be a statement of the forcing language appropriate for $\mathcal{P}_\alpha$. Then there is a direct extension p^* of p deciding σ .*

Proof: Suppose otherwise. Let $p = \langle p_\gamma \mid \gamma \in g \rangle$ and β be the minimal element of g . We assume that $g \neq \emptyset$, otherwise any extension of it is direct.

Let G be a generic subset of $\mathcal{P}_{\beta+1}$, so that $p \restriction \beta + 1 \in G$. We shall mean by $p \setminus (\beta + 1) = \langle p_\gamma \mid \gamma \in g \setminus (\beta + 1) \rangle$ the interpretation of it in $V[G]$, i.e. an element of the forcing $\mathcal{P}_\alpha/G$. Define now $p^* \in \mathcal{P}_\alpha/G$. If there exists some $q \in \mathcal{P}_\alpha/G$ a direct extension of $p \setminus (\beta + 1)$ deciding σ , then set p^* to some such q . Otherwise, set $p^* = p \setminus (\beta + 1)$. Let $p^* = \langle p_\gamma^* \mid \gamma \in g^* \rangle$. Then $g^* \in V[G]$. But, since $\mathcal{P}_{\beta+1}$ is of small cardinality (see (3) of the definition of $\mathcal{P}_\alpha$) g^* can be easily replaced by a set in V . Suppose that is already the case. Then $p \restriction (\beta + 1) \cup p^*$ will be a condition in $\mathcal{P}_\alpha$ and it will be a direct extension of p .

Claim 1.4.1. There is p_β^* such that $\phi \Vdash_{\mathcal{P}_\beta} “p_\beta^* \geq p_\beta^*”$ and $\langle p_\beta^* \rangle \Vdash_{\mathcal{P}_{\beta+1}}$

$$“p^* = p \setminus (\beta + 1)”$$

Proof: Let $G \subseteq \mathcal{P}_\beta$ be generic. Work in $V[G]$. Let p_β^* be a direct extension of p_β (in Q_β) deciding the statement “ $p^* = p \setminus (\beta + 1)$ ”. Suppose for a moment that the decision is

negative. Then $p_\beta^* \Vdash_{Q_\beta} "p^* \text{ is a direct extension of } p \setminus (\beta + 1) \text{ deciding } \sigma"$, by the choice of p^* . Now pick a direct extension p_β^{**} of p_β^* so that

$$p_\beta^{**} \Vdash_{Q_\beta} "p^* \Vdash^i \sigma",$$

where ${}^\circ \sigma = \sigma$ and ${}^1 \sigma = {}^\neg \sigma$. Pick some $r \in G$ so that

$$r \cap \langle p_\beta^{**} \rangle \Vdash_{\mathcal{P}_{\beta+1}} "p^* \Vdash^i \sigma".$$

Then $r \cap \langle p_\beta^{**} \rangle \cap p^*$ will be a direct extension of p forcing ${}^i \sigma$. Which contradicts our assumption. $\square$ of the claim.

Now we replace p_β in p by p_β^* . Denote the resulting condition by $p(\{\beta\})$. Let β_1 be the second element of g . We proceed as above replacing β by β_1 and p by $p(\{\beta\})$. This will define p_{β_1} . Then set $p(\{\beta_0, \beta_1\}) = \langle p_\beta^*, p_{\beta_1}^* \rangle \cap \langle p_\gamma \mid \gamma \in g \setminus \{\beta, \beta_1\} \rangle$. Continue in the same fashion. Finally, after going through all the elements of g we will obtain a condition $p(g) = \langle p_\gamma^* \mid \gamma \in g \rangle$ which is a direct extension of p . Now let $q \geq p(g)$ be a condition deciding σ . By the definition of extension, there is a maximal $\gamma \in g$ such that $q \restriction \gamma \Vdash_{\mathcal{P}_\gamma} "q_\gamma \text{ is not a direct extension of } p_\gamma"$. But this will contradict the choice of $p(g \cap (\gamma + 1))$, since above γ , $q \setminus \gamma$ will be a direct extension of $p \setminus \gamma$ deciding σ . Contradiction. $\square$

Now let us define the iteration needed for the proof of 1.2.

Let $\alpha < \kappa$ be an ordinal with $o(\alpha) = \alpha$. We like to define Q_α . But first, let us consider the forcing $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$, where $\beta < \alpha$. Define a $*$ -ordering on $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ by setting $p = \langle b, a \rangle \leq^* p' = \langle b', a' \rangle$ if $p = p'$ or (i) $p \leq p'$ and (ii) $b \leq^* b'$ in $\mathcal{P}(\alpha, \beta)$.

Lemma 1.4. *The forcing $\langle \mathcal{P}(\alpha, \beta) * P[E \cap \alpha], \leq, \leq^* \rangle$ satisfies the Prikry condition.*

The proof follows easily from Fact 1.

Lemma 1.5.0. *Every condition in $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ can be extended to a condition of the form $\langle b, \check{c} \rangle$.*

It follows by Fact 1 since $\mathcal{P}(\alpha, \beta)$ does not add new bounded subsets to α .

Let $D_{\alpha, \beta} = \{ \langle b, a \rangle \in \mathcal{P}(\alpha, \beta) * P[E \cap \alpha] \mid b \Vdash_{\mathcal{P}(\alpha, \beta)} " \sup a \in \check{B} "$, where $\check{B}$ is the canonical name of the closed cofinal in α sequence added by $\mathcal{P}(\alpha, \beta)$.

Lemma 1.6. For every $p \in \mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ there exists $q \in D_{\alpha, \beta}$ such that $q \geq^* p$.

Proof: Obviously, since $b \Vdash_{\mathcal{P}(\alpha, \beta)} \sup \check{a} < \alpha$, where $p = \langle b, \check{a} \rangle$. $\square$

Lemma 1.7. The order $\leq^*$ is β -closed over $D_{\alpha, \beta}$ and hence $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ does not add new bounded subsets of β .

Remark. Notice that all cardinals $\leq \alpha$ are collapsed to β , since the forcing with $P[E \cap \alpha]$ over $V^{\mathcal{P}(\alpha, \beta)}$ is isomorphic to the Levy collapse $\text{Col}(\beta, \alpha)$.

Proof: Let $\langle \check{b}_i, \check{a}_i \mid i < \tau < \beta \rangle$ be a $\leq^*$ -increasing sequence of elements of $D_{\alpha, \beta}$. Since $\langle \mathcal{P}(\alpha, \beta), \leq^* \rangle$ is α -closed, there is $b \in \mathcal{P}(\alpha, \beta)$ such that $b \geq^* \check{b}_i$ for all $i < \tau$. Let $\check{a}$ be a name of the union of $\check{a}_i$'s. Since for every $i < \tau$ $b \Vdash''_{\mathcal{P}(\alpha, \beta)} \sup(\check{a}_i) \in \check{B}''$ where $\check{B}$ is a canonical name of the generic closed cofinal in α sequence, $b \Vdash_{\mathcal{P}(\alpha, \beta)} \text{"}\sup \check{a} \in \check{B}\text{"}$ and in particular $b \Vdash_{\mathcal{P}(\alpha, \beta)} \text{"}\sup \check{a} \text{ is regular in } V\text{"}$. Hence $\langle b, \check{a} \rangle$ is a condition, it belongs to $D_{\alpha, \beta}$ and it is $\leq^*$ stronger than each $\langle \check{b}_i, \check{a}_i \rangle$ ($i < \tau$). $\square$

Definition 1.8.

- (1) $Q_\alpha = P[E \cap \alpha] \cup \bigcup_{\beta < \alpha} (\mathcal{P}(\alpha, \beta) * P[E \cap \alpha])$.
- (2) The ordering of Q_α :
- (2a) The ordering of $P[E \cap \alpha]$ and of $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ ($\beta < \alpha$) is the usual one
- (2b) Let $c \in P[E \cap \alpha]$ and $\langle b, \check{a} \rangle \in \mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$, for some $\beta < \alpha$. Define

$$\begin{aligned} \langle b, \check{a} \rangle \geq c &\text{ iff } b \Vdash_{\check{a} \geq_{P[E \cap \alpha]} \check{c}} \\ c \geq \langle b, \check{a} \rangle &\text{ iff } b \Vdash_{\check{c} \geq_{P[E \cap \alpha]} \check{a}} \end{aligned}$$

- (2c) Let $\langle b_i, \check{a}_i \rangle \in \mathcal{P}(\alpha, \beta_i) * P[E \cap \alpha]$ where $i = 0, 1, \beta_0 \neq \beta_1 < \alpha$. Define $\langle b_0, \check{a}_0 \rangle \leq \langle b_1, \check{a}_1 \rangle$ iff there is $c \in P[E \cap \alpha]$ such that $\langle b_0, \check{a}_0 \rangle \leq c \leq \langle b_1, \check{a}_1 \rangle$.

We also define $\ast$ -ordering of Q_α .

Definition 1.9. Let $p, q \in Q_\alpha$. Set $p \leq^* q$ iff $p = q$ or for some $\beta < \alpha$ $p = \langle b_1, \check{a}_1 \rangle$, $q = \langle b_2, \check{a}_2 \rangle \in \mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ and (i) $p \leq q$ (ii) $b_1 \leq^* b_2$ in $\mathcal{P}(\alpha, \beta)$.

Lemma 10. (1) $P[E \cap \alpha]$ is dense in Q_α .

(2) The forcing $\langle Q_\alpha, \leq \rangle$ is equivalent to $\langle P[E \cap \alpha], \leq \rangle$ and hence preserves cofinalities of cardinals.

Proof: $P[E \cap \alpha]$ is dense in Q_α by the definition of the order. Also conditions incompatible in $P[E \cap \alpha]$ remaining so in Q_α . $\square$

Lemma 11. *Let $\beta < \alpha$. Suppose that $\langle b, a \rangle \in \mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ and σ is a statement of the forcing language of Q_α . Then there is a direct extension (in Q_α) of $\langle b, a \rangle$ deciding σ in the forcing with Q_α .*

Remark. The lemma actually shows the Prikry condition above $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ part of Q_α .

The lemma will follow from the following statement.

Lemma 12. *For every $\beta < \alpha$ Q_α is a projection of $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$.*

Proof: It is enough to project a dense subset of $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$ onto a dense subset of Q_α . Consider $D \subseteq \mathcal{P}(\alpha, \beta) \times P[E \cap \alpha]$, which is dense in $\mathcal{P}(\alpha, \beta) * P[E \cap \alpha]$. There is such D since $\mathcal{P}(\alpha, \beta)$ does not add new bounded subsets of α . For $\langle b, a \rangle \in D$ set $\pi(\langle b, a \rangle) = a$. By Lemma 10, $rng \pi = P[E \cap \alpha]$ is dense in Q_α . It is trivial that π is a projection map. $\square$

We are ready now to define the iteration. The definition will be as above only β 's for Q_α will be picked generically more carefully. This is needed for cardinals preservation.

Let A and A^ℓ be as above. For every $\alpha \in A^\ell$ we define $\mathcal{P}_\alpha$ by induction.

Definition 13. A forcing notion $\mathcal{P}_\alpha$ consists of all elements p of the form $\langle p_\gamma \mid \gamma \in g \rangle$ where

- (1) g is a subset of $\alpha \cap A$;
- (2) g has an Easton support;
- (3) for every $\gamma \in \text{dom } g$ $p \restriction \gamma = \langle p_\beta \mid \beta < \gamma \rangle \in \mathcal{P}_\gamma$ and $p \restriction \gamma \Vdash''_{\mathcal{P}_\gamma} p_\gamma$ is a condition in either the forcing Q_γ , if $o(\gamma) = \gamma$ or in the forcing $\mathcal{P}(\gamma, o(\gamma))$, if $o(\gamma) < \gamma$;
- (4) for every $\tau \leq \alpha$ the set $\{\gamma < \alpha \mid p_\gamma \in P_\tau[E \cap \gamma]\}$ or $\gamma \geq \tau$ and for some $\beta < \tau$ $p_\gamma \in \mathcal{P}_\tau(\gamma, \beta) * P_\tau[E \cap \gamma]\}$ is finite.

The ordering on $\mathcal{P}_\alpha$ is defined without changes.

The definition and Lemma 1.10 insures that for every γ with $o(\gamma) = \gamma$ the actual forcing used over γ is $P[E \cap \gamma]$. But in every separate condition $p \in \mathcal{P}_\alpha$ only finitely many γ 's with $p_\gamma \in P[E \cap \gamma]$ are allowed (the condition (4)). The reason for this is to insure that

the iteration preserves cardinals. Intuitively, finite iteration of forcings $P[E \cap \gamma]$'s does no harm. In order to do infinite iterations (even of the length ω), we like to have in advance some information about closed pieces of E . Forcings $\mathcal{P}(\gamma, \beta)$'s are actually used for this purpose. Namely canonical generic sequences produced by such forcings.

Lemma 14. $\langle \mathcal{P}_\alpha, \leq, \leq^* \rangle$ satisfies the Prikry condition.

The proof repeats the proof of Lemma 13. The additional condition (4) has no effect on it.

Lemma 15. Let $\gamma < \alpha$. Suppose that G is a generic subset of $\mathcal{P}_{\gamma+1}$. Then the forcing $\mathcal{P}_\alpha/G$ does not add new subsets of γ .

Proof: Suppose that some $p \Vdash a \subseteq \check{\gamma}$. It is enough to show the following.

Claim 1.15.1. There are $b \in V[G]$ and $\bar{p} \geq p$ such that $\bar{p} \Vdash a = \check{b}$.

Remark. $\bar{p}$ is not required to be a direct extension of p . The reason for this is the finite set of γ 's in p satisfying (4).

Proof: Using Lemma 1.14, we define by induction a $*$ -increasing sequence $\langle p_i \mid i \leq \gamma \rangle$ of extensions of p so that for each i

- (a) p_i decides the statement “ $i \in a$ ”.
- (b) if for some δ with $o(\delta) = \delta$ $\delta \in g_i$, then $p_i \restriction \delta \Vdash_{\mathcal{P}_\delta}$ “ $p_{i\delta} \in D_{\delta, \beta}$ for some $\beta > \gamma$ ” where $p_i = \langle p_{i\nu} \mid \nu \in g_i \rangle$,
- (c) if $j > i$, $j \leq \gamma$, then for every $\delta \in g_j \setminus g_i$ with $o(\delta) = \delta$

$$p_j \restriction \delta \Vdash “p_{j\delta} \in \mathcal{P}(\delta, \beta) * P[E \cap \delta]”$$

for some β which is above $\sup(g_i \cap \delta)$.

First we extend p to a condition p' satisfying (b). By (4) of Definition 1.13 it is always possible. But p' need not be a direct extension of p . Now, by Lemma 1.9, find $p'' \geq^* p'$ deciding “ $0 \in a$ ”. Let p_0 be an extension of p'' obtained as follows. We replace each p_j in p'' for δ with $0(\delta) = \delta$ which is not in p' by a stronger condition in $\mathcal{P}(\delta, \beta) * P[E \cap \delta]$ where β is picked to be above every coordinate of p' below δ . By (3), (4) of Definition 1.13,

only finitely many coordinates δ in p'' should be fixed this way. So p_0 will be a condition stronger than p'' and a direct extension of p' deciding “ $\check{0} \in a$ ”.

We continue by induction. On successor stages we proceed as above. Suppose now that $i \leq \gamma$ is a limit ordinal and a sequence $\langle p_\rho \mid \rho < i \rangle$ is defined and satisfies the conditions (a) - (c) above. Let us argue that there is p' a direct extension of all of p_ρ 's ($\rho < i$). Let p' be obtained by taking direct extensions in each coordinate separately. This is possible by (b). It is enough to show that such an obtained p' is a condition. The only problematic point is (4) of Definition 1.13. By (b), $\{\gamma < \alpha \mid p'_\gamma \in P[E \cap \gamma]\}$ is empty. So it remains to show that for every $\tau < \alpha$ the set $\{\gamma < \alpha \mid \tau \leq \gamma, \text{ for some } \beta < \tau \ p'_\gamma \in \mathcal{P}(\gamma, \beta) * P[E \cap \gamma]\}$ is finite. Suppose otherwise. Let $\tau, \tau \leq \gamma_0 < \gamma_1 < \dots < \gamma_n < \dots < \alpha$ be witnessing this. Then for each $n < \omega$ there is $\beta_n < \tau$ such that $p'_{\gamma_n} \in \mathcal{P}(\gamma_n, \beta_n) * P[E \cap \gamma_n]$. For each $n < \omega$ let $i_n < i$ be the least such that the coordinate γ_n appears in p_{i_n} . Shrinking the set of indexes if necessary, we assume that the sequence $\langle i_n \mid n < \omega \rangle$ is strictly increasing. But this is impossible by (c). Since $\gamma_1 > \gamma_0$, γ_1 is in p_{i_1} and not in p_{i_0} , γ_0 is in p_{i_0} but $p'_{i_1 \gamma_1} \in \mathcal{P}(\gamma_1, \beta_1) * P[E \cap \gamma_1]$ where $\beta_1 < \tau < \gamma_0$. Contradiction. So p' is a condition. Now we continue as in successor stages. $\square$ of the claim.

$\square$

Lemma 1.6. $\mathcal{P}_\kappa$ satisfies κ -c.c. and preserves the cardinals.

Follows from Definition 1.13 and Lemma 1.15.

Let G_κ be a generic subset of $\mathcal{P}_\kappa$. Force with $P[E]$ over $V[G_\kappa]$. Let C be a generic club.

Lemma 1.17. κ is a measurable cardinal in $V[G_\kappa, C]$.

Proof: Let U be the measure in V concentrating over $\{\alpha < \kappa \mid o(\alpha) = \alpha\}$. Denote by $i_u : V \rightarrow N_u \simeq \text{Ult}(V, U)$ the corresponding elementary embedding. Since $\mathcal{P}_\kappa$ satisfies κ -c.c. and using Claim 1.15.1 it is routine to extend i_u (in $V[G_\kappa, C]$) to an embedding

$$i : V[G_\kappa] \longrightarrow N[G_{i_u(\kappa)}]$$

where N is an iterated ultrapower of N_u moving only ordinals in the interval $(\kappa^+, i_u(\kappa))$.

We refer to [Gi1,3] or [Gi-Sh] for such arguments.

For the benefit of the reader, let us provide more details on what is going on.

Working in $V[G_\kappa, C]$ we define a normal $V[G_\kappa]$ -ultrafilter over κ (i.e. a normal ultrafilter over $\mathcal{P}^{V[G_\kappa]}(\kappa)$). Proceed as follows: pick in V an enumeration $\langle \mathcal{D}_\alpha \mid \alpha < \kappa^+ \rangle$ such that in N_U

$\emptyset \Vdash_{\mathcal{P}_{i_u(\kappa)}/G_\kappa * C} \text{“}\mathcal{D}_\alpha \text{ is a } * \text{-dense open subset of } i_u(\mathcal{P}_\kappa)/\mathcal{P}_{\kappa+1}\text{”}$ and if for some $\mathcal{D} \Vdash_{\mathcal{P}_{i_u(\kappa)}/G_\kappa * C} \text{“}\mathcal{D} \text{ is a } * \text{-dense open subset of } i_u(\mathcal{P}_\kappa)/\mathcal{P}_{\kappa+1}\text{”}$ then for some $\alpha < \kappa^+$ $\emptyset \Vdash \mathcal{D} \subseteq \mathcal{D}_\alpha$.

Then define a master condition sequence $\langle r_\alpha \mid \alpha < \kappa^+ \rangle \in V[G_\kappa, C]$ such that

- (a) $r_\alpha \leq^* r_\beta$ for every $\alpha \leq \beta < \kappa^+$
- (b) $r_\alpha \in D_\alpha[G_\kappa, C]$ for every $\alpha < \kappa^+$.
- (c) $\langle r_\alpha \mid \alpha < \beta \rangle \in N_u[G_\kappa, C]$ for every $\beta < \kappa^+$.

Using $\langle r_\alpha \mid \alpha < \kappa^+ \rangle$ we define now $U^* \subseteq U$ by setting $X \in U^*$ iff for some $p \in G_\kappa * C$, $\alpha < \kappa^+$, a $\mathcal{P}_\kappa$ -name $\check{X}$ of X , in N_u

$$\langle p, r_\alpha \rangle \Vdash_{i_u(\mathcal{P}_\kappa)} \text{“}\check{X} \in X\text{”}.$$

It is easy to check that U^* is a normal $V[G_\kappa]$ -ultrafilter. Moreover, since $P[E]$ does not add new sequences of the length less than κ to $V[G_\kappa]$, U^* is κ -complete in $V[G_\kappa, C]$. Hence $Ult(V[G_\kappa], U^*)$ is well-founded. Let N^* be its transitive collapse and $i : V[G_\kappa] \longrightarrow N^*$ be the corresponding elementary embedding. By elementarity N^* is of the form $N[G_{i(\kappa)}]$ where $G_{i(\kappa)}$ is N -generic subset of $\mathcal{P}_{i(\kappa)}$. Examining the structure of N it is possible to show that it is in fact an iterated ultrapower of N_u moving only ordinals in the interval $(\kappa^+, i_u(\kappa))$. But this is not needed for further argument.

Since $P[E]$ does not add new $< \kappa$ -sequences, $N[G_{i(\kappa)}]$ is closed under $< \kappa$ -sequences of its elements. But really more is true:

Claim 1.17.1. ${}^\kappa N[G_{i(\kappa)}] \subseteq N[G_{i(\kappa)}]$.

Proof: Let $\langle t_\alpha \mid \alpha < \kappa \rangle$ be a sequence of elements of $N[G_{i(\kappa)}]$. Without loss of generality, we may assume that each t_α is an ordinal. Let $\check{t}$ be a canonical $\mathcal{P}_\kappa * P[E]$ name of this sequence. Then for each α $\check{t}(\alpha)$ is a set of cardinality at most κ consisting of pairs $\langle p, \check{\delta} \rangle$ where $p \in \mathcal{P}_\kappa * P[E]$ and δ is an ordinal. For every $\alpha, \gamma < \kappa$ such that $o(\gamma) = \gamma$ let $\check{t}(\alpha) \upharpoonright \gamma$

denotes the set of γ first pairs $\langle p, \check{f}_\delta(\gamma) \rangle$ in $\check{t}(\alpha)$ such that $p \in \mathcal{P}_\gamma * P[E \cap \gamma]$ where f_δ represents δ in N_u . Set $\check{t} \restriction \gamma = \langle \check{t}(\alpha) \restriction \gamma \mid \alpha < \gamma \rangle$. Then, $i_u(\langle \check{t} \restriction \gamma \mid \gamma < \kappa \rangle)(\kappa) = \check{t}$. Now define a function $g \in V[G_\kappa]$ representing $\langle t_\alpha \mid \alpha < \kappa \rangle$ in $N[i_u(\mathcal{P}_\kappa)]$. Set $g(\gamma) = \check{t} \restriction \gamma[G_\gamma * C_\gamma]$, where $G_\gamma * C_\gamma = G_{\gamma+1} = G_\kappa \cap \mathcal{P}_\gamma * P[E \cap \gamma]$. $\square$

Now, in $V[G_\kappa]$ there is a Rudin-Kiesler increasing commutative sequence of ultrafilters $\langle U_\alpha \mid \alpha < \kappa \rangle$ over κ . Thus for each $\alpha < \kappa$ the measure concentrating on $\{\beta < \kappa \mid o(\beta) = \alpha\}$ in V extends to U_α in $V[G_\kappa]$. Actually, such extensions are used to define $\mathcal{P}(\gamma, \delta)$'s. Then, in $N[G_{i(\kappa)}]$ we will have such a sequence over $i(\kappa)$ of the length $i(\kappa)$. Form the direct limit. Let $k : N[G_{i(\kappa)}] \rightarrow M[G_\lambda]$ be the corresponding embedding, where $\lambda = k(i(\kappa))$. Let $j = k \circ i : V[G_\kappa] \rightarrow M[G_\lambda]$. Notice that since the length of the sequence used to form the direct limit is $i(\kappa)$ and $\text{cf}(i(\kappa)) = \kappa^+$, ${}^\kappa M[G_\lambda] \subseteq M[G_\lambda]$. An additional point here is that λ is a limit of critical points of embeddings used in the direct limit. They are singular cardinals in $M[G_\lambda]$ but are regular in M . Since M is just an iterated ultrapower of $\mathcal{K}$, by [Mi2]. Hence, in $V[G_\kappa, C]$, $j(E)$ contains a club. Now we use this in the standard fashion to diagonalize over κ^+ dense subset of $(P[j(E)])^{M[G_\lambda]}$. It will produce a club $C^* \supseteq C$ which is $M[G_\lambda]$ -generic. Notice that by Lemma 1.10(2), Q_κ is equivalent to $P[E \cap \kappa]$. So C is Q_κ generic. So we obtain $j \subseteq j^* : V[G_\kappa, C] \rightarrow M[G_\lambda, C^*]$. Hence κ is measurable in $V[G_\kappa, C]$. $\square$

2. On the strength of the existence of a club of “regulars”

In this section we will show that the hypothesis used in Theorems 1.1 are the best possible and those of 1.2 are close to this.

The next theorem is basically due to Mitchell.

Theorem 2.1. *Suppose that κ is an inaccessible and the set $E = \{\alpha < \kappa \mid \alpha \text{ is regular in } \mathcal{K}\}$ contains a club. Then for every $\delta < \kappa$ the set $\{\alpha < \kappa \mid o^{\mathcal{K}}(\alpha) \geq \delta\}$ is stationary.*

Proof: Let $\delta < \kappa$. We show that $\{\alpha < \kappa \mid o^{\mathcal{K}}(\alpha) \geq \delta\} = A_\delta$ is stationary. Let C be a club contained in E . Choose some $\alpha \in C$ of cofinality δ . Then $\text{cf}^{\mathcal{K}} \alpha = \alpha$ since $\alpha \in E$. So its cofinality changed to δ . Then by Mitchell [Mi2], $o^{\mathcal{K}}(\alpha) \geq \delta$. Hence $C \cap A_\delta \neq \emptyset$. $\square$

Theorem 2.2. *Let κ be a measurable cardinal and a set $E = \{\alpha < \kappa \mid \alpha \text{ is regular in } \mathcal{K}\}$ contain a club. Then $o^\mathcal{K}(\kappa) \geq \kappa$.*

Proof: First note that by Mitchell [Mi3], $\kappa^+ = (\kappa^+)^{\mathcal{K}}$. Let U be a normal measure over κ . Consider its elementary embedding $j_U : V \rightarrow N \simeq \text{Ult}(V, U)$. By Mitchell [Mi3], $j_U \upharpoonright \mathcal{K} = i$ is an iterated ultrapower of $\mathcal{K}$ by its measures. Suppose that $o^\mathcal{K}(\kappa) < \kappa$. Then $i(\kappa)$ is not a limit point of iteration. Since ${}^\kappa N \subseteq N$, cf $i(\kappa) \geq \kappa^+$. That is, there is a last measure in which the ultrapower reached $i(\kappa)$, (or more precisely, the image of its critical point). Denote this critical point by λ . Then for every $\alpha < i(\kappa)$ there is $f : [\kappa]^n \rightarrow \kappa$ ($n < \omega$) in $\mathcal{K}$ and $\kappa \leq \beta_1, \dots, \beta_n \leq \lambda$ such that $i(f)(\beta_1, \dots, \beta_n) = \alpha$.

Using coding of n -tuples, we can replace n -placed functions by a 1-placed ones. Then, for every $\alpha < i(\kappa)$ (or $g : \kappa \rightarrow \kappa$) there will be $\beta \leq \lambda$ and $f : \kappa \rightarrow \kappa$ in $\mathcal{K}$ such that $i(f)(\beta) = \alpha$ (or $i(f)(\beta) = j_U(g)(\kappa)$).

Claim 1. *For every $\alpha < i(\kappa)$ there is $f \in {}^\kappa \kappa \cap \mathcal{K}$ such that $\alpha \leq i(f)(\lambda)$.*

Proof: Let $\alpha < i(\kappa)$ and $g \in {}^\kappa([\kappa]^n) \cap \mathcal{K}$ be such that $i(g)(\beta_1, \dots, \beta_{n-1}, \lambda) = \alpha$, where $\beta_1 < \dots < \beta_{n-1} < \lambda$. Define $f \in {}^\kappa \kappa \cap \mathcal{K}$ as follows:

$$f(\nu) = \sup\{g(\nu_1, \dots, \nu_{k-1}, \nu) \mid \nu_1 < \dots < \nu_{n-1} < \nu\} .$$

□

Claim 2. *The set*

$$A = \{\beta < i(\kappa) \mid \exists f \in {}^\kappa \kappa \cap \mathcal{K}, \quad i(f)(\lambda) = \beta\}$$

is $< \kappa$ -closed.

Proof: First suppose that the embedding i is definable in $\mathcal{K}$. Let $\{\beta_\nu \mid \nu < \rho\}$ be a subset of A for some $\rho < \kappa$. Denote by β the $\sup\{\beta_\nu \mid \nu < \rho\}$. Since there is no measurable cardinals of $\mathcal{K}$ between κ and 2^κ , there will be a set $B \in \mathcal{K}$, $|B| \leq \kappa$ covering $\{\beta_\nu \mid \nu < \rho\}$, by Mitchell [Mi2]. Notice that since $\kappa^+ = (\kappa^+)^{\mathcal{K}}$, $|B|^\mathcal{K} \leq \kappa$. Using the regularity of κ , we can find $B^* \subseteq B \cap A$, $B^* \in \mathcal{K}$ of cardinality $< \kappa$ cofinal in β .

Now it is obvious that for a function $f \in {}^\kappa \kappa \cap \mathcal{K}$, $i(f)(\lambda) = \beta$. So $\beta \in A$.

Let us now deal with the general case, i.e. i is not necessarily definable in $\mathcal{K}$. We replace i by a definable in $\mathcal{K}$ iteration i^* . Proceed as follows. Iterate $\mathcal{K}$ using every measure over κ as well as the new ones appearing in the process $(2^\kappa)^+$ -many times. Let i^* be such an iteration. Obviously, there is an embedding $k : i(\mathcal{K}) \rightarrow i^*(\mathcal{K})$ so that the following diagram is commutative:

$$\begin{array}{ccc} & & i(\mathcal{K}) \\ & \nearrow i & \downarrow k \\ \mathcal{K} & & i^*(\mathcal{K}) \\ & \searrow i^* & \end{array} \quad .$$

Notice that $k \restriction \kappa^+ = id$. So, $k(A) = \{\beta < i^*(\kappa) \mid \exists f \in {}^\kappa \kappa \cap \mathcal{K}, i^*(f)(k(\lambda)) = \beta\} = k''(A)$.

Now, the argument used above for i , A works for i^* , $k(A)$. $\square$

Let us now change the cofinality of κ to ω by the Prikry forcing with U . Let $\langle \kappa_n \mid n < \omega \rangle$ be the Prikry sequence and $\langle \lambda_n \mid n < \omega \rangle$ the sequence corresponding to λ , i.e. $\langle f(\kappa_n) \mid n < \omega \rangle$ for $f \in {}^\kappa \kappa$ representing λ in the ultrapower by U .

Choose an elementary submodel M , of large enough portion of the universe $|M| < \kappa$, ${}^\omega M \subseteq M$ containing all relevant information. Let $\alpha = \sup(M \cap i(\kappa))$ and $\alpha_n = \sup(M \cap \kappa_{n+1})$ for $n < \omega$. Notice that $\text{cf } \alpha < \kappa$, since $|M| < \kappa$ and $\text{cf } \alpha > \omega$, since ${}^\omega M \subseteq M$.

Since $A = \{\beta < i(\kappa) \mid \exists f \in {}^\kappa \kappa \cap \mathcal{K}, i(f)(\lambda) = \beta\}$ is $< \kappa$ -closed and unbounded in $i(\kappa)$ in V , it contains its limit points of cofinality $\delta, \omega < \delta < \kappa$, in $V[\langle \kappa_n \mid n < \omega \rangle]$. Hence, $\alpha \in A$, since $A \in M$ and $\omega < \text{cf } \alpha < \kappa$. Let $f_\alpha \in {}^\kappa \kappa \cap \mathcal{K}$ be such that $i(f_\alpha)(\lambda) = \alpha$.

Claim 3. *For all but finitely many n 's, $f_\alpha(\lambda_n) = \alpha_n$.*

Proof: Let g_ν be a function in V such that $\nu = i(g_\nu)(\kappa)$. Then it is enough to show that $g_\alpha(\kappa_n) = \alpha_n$ for almost all n , for then

$$\begin{aligned} i(f_\alpha)(\lambda) = \alpha &\iff \{\nu : f_\alpha(g_\lambda(\nu)) = g_\alpha(\nu)\} \in U \\ &\iff f_\alpha(g_\lambda(\kappa_n)) = g_\alpha(\kappa_n) \text{ for sufficiently large } n \\ &\iff f_\alpha(\lambda_n) = \alpha_n \text{ for sufficiently large } n . \end{aligned}$$

Now we show that $g_\alpha(\kappa_n) = \alpha_n$ for almost every n . Let $g \in M \cap \prod_{n < \omega} \kappa_{n+1}$. By standard argument on Prikry forcing used inside M , we can find $\beta \in M \cap \alpha$ such that $g_\beta(\kappa_n) > g(n)$ for all but finitely many n 's. So $\alpha_n \leq g_\alpha(\kappa_n)$. Now $\omega < \text{cf}(\alpha) < \kappa$ in V since $\omega < \text{cf}(\alpha) < \kappa$ in $V[(\kappa_n : n \in \omega)]$. Choose an increasing sequence $(\xi_\nu : \nu < \rho) \in V$ which is cofinal in α , for some $\rho < \kappa$. Then

$$\left\{ \eta : g_\alpha(\eta) = \sup_{\nu < \rho} g_{\xi_\nu}(\eta) \right\} \in U .$$

If $\alpha_n < g_\alpha(\kappa_n)$ for finitely many n then there is $\nu < \rho$ such that $\alpha_n < g_{\xi_\nu}(\kappa_n)$ for infinitely many n (using the fact that $\text{cf}(\alpha) > \omega$). This is impossible, since if we pick $\xi \in M$ with $\xi_\nu < \xi < \alpha$ then $g_{\xi_\nu}(\kappa_n) < g_\xi(\kappa_n) < \alpha_n$ for almost all n . $\square$

Let us now use the club $E \in V$ consisting of regular in $\mathcal{K}$ cardinals. For every $n < \omega$, $E \cap \kappa_{n+1}$ will be such a club in κ_{n+1} . Hence the same is true in M . So, $\alpha_n = M \cap \kappa_{n+1} \in E$ for every $n < \omega$. We argue that this is impossible.

Briefly we apply Mitchell's analysis of M , see [Mi2]. It implies that for all but finitely many n 's, α_n is a limit of indiscernibles. Since κ_{n+1} is indiscernible for κ , there will a club subset of $\alpha_n \cap E$ consisting of indiscernibles for at least κ . Select a sequence $\langle c_n \mid n < \omega \rangle$ of such indiscernibles so that $\lambda_n < c_n < \alpha_n$. Then $\langle c_n \mid n < \omega \rangle \in M$ and by Claim 3, $f_\alpha(\lambda_n) > c_n$ ($n < \omega$), which is impossible. A contradiction.

Let us provide more details.

We apply the technique of the second section of [Mi2]. We will use M as the set N of that paper, i.e. we are covering the set M .

Write $C(\alpha) = \bigcup_\beta \mathcal{C}(\alpha, \beta)$ and $C = \bigcup_\alpha C(\alpha)$. It follows from what is given there that there are $h^M \in \mathcal{K}$ and $\xi < \kappa$ such that:

- (1)(a) $\forall \gamma \in M \cap \kappa \ \gamma \in h^{M''}(\xi \cup (C \cap \gamma + 1))$.
- (b) If for $c \in C$ we write $\tau(c)$ for the largest ordinal τ such that $c \in \mathcal{C}(\tau)$ then $\tau(c)$ always exists and is in $h^{M''}(\xi \cup (C \cap c))$.

Since α_n is regular in $\mathcal{K}$, it follows that C is unbounded in α_n and $\kappa_n \in C$ for sufficiently large n . Since $(\kappa_n : n \in \omega)$ is a Prikry sequence it follows that $\tau(\kappa_n) = \kappa$ for sufficiently large n . Also it follows that for c in a closed unbounded subset of $\alpha_n \cap M$ we have $\tau(c) \not\leq \alpha_n$. Since $\tau(c) \in M$ it follows that $\tau(c) > \alpha_n$.

An additional fact from [Mi2] that we are using:

(2) For every $f \in \mathcal{K}$ there is $\rho < \kappa$ such that if $c > \rho$ then $(f''c) \cap \tau(c) \subset c$.

However we have we have $f_\alpha(\lambda_n) = \alpha_n$ for all sufficiently large n , and this is impossible because for all sufficiently large n there are c_n such that $\lambda_n < c_n < \alpha_n$ and $\tau(c_n) > \alpha$. $\square$

We do not know if it is possible to improve the above to $o(\kappa) = \kappa + 1$.

Question. Is $o(\kappa) = \kappa$ enough for a model with a measurable containing a club of former regular cardinals?

References

- [Av-Sh] U. Avraham and S. Shelah, Forcing closed unbounded sets, *J. Sym. Logic* 48 (1983), 643-648.
- [Gi1] M. Gitik, Changing cofinalities and the nonstationary ideal, *Israel Journal of Math.* 56(3) (1986), 280-314.
- [Gi2] M. Gitik, On measurable cardinals violating GCH, *Ann. of Pure and Appl. Logic* 63 (1993), 227-240.
- [Gi3] M. Gitik, On the Mitchell and Rudin-Kiesler orderings of ultrafilters, *Ann. of Pure and appl. Logic* 39 (1988), 175-197.
- [Gi-Sh] M. Gitik and S. Shelah, On Certain Indestructibility of Strong Cardinals and a Question of Hajnal, *Arch. Math. Logic* 28, (1989) 35-42.
- [Mi1] W. Mitchell, How weak is a closed unbounded ultrafilter? In *Logic Coll.* 80, North Holland 1982, pp. 209-231.
- [Mi2] W. Mitchell, Applications of the Core Model for sequences of measures, *Trans. Amer. Math. Soc.* 299 (1987), 41-58.
- [Mi3] W. Mitchell, The Core Model for sequences of measures I, *Math. Proc. Cambridge Phil. Soc.* 95 (1984), 229-260.